

HUMAN BAD POLYPEPTIDES, ENCODING NUCLEIC
ACIDS AND METHODS OF USE

Throughout this application various publications are referenced within parentheses. The disclosures of these publications in their entireties are hereby incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains.

BACKGROUND OF THE INVENTION

- 10 The present invention relates generally to apoptosis or, programmed cell death, and more particularly, nucleic acids encoding human Bad polypeptides which can be used to modulate apoptosis for the therapeutic treatment of human diseases.
- 15 Apoptosis is a normal physiological process of cell death that plays a critical role in the regulation of tissue homeostasis by ensuring that the rate of new cell accumulation produced by cell division is offset by a commensurate rate of cell loss due to death. It has
- 20 now become clear that disturbances in apoptosis, also referred to as physiological cell death or programmed cell death, that prevent or delay normal cell turnover can be just as important to the pathogenesis of diseases as are known abnormalities in the regulation of
- 25 proliferation and the cell cycle. Like cell division, apoptosis is similarly regulated under normal circumstances through complex interaction of gene products that either promote or inhibit cell death.

The stimuli which regulate the function of these apoptotic gene products include both extracellular and intracellular signals. Either the presence or the absence of a particular stimuli can be sufficient to evoke a positive or negative apoptotic signal. For example, physiological stimuli that prevent or inhibit apoptosis include, for example, growth factors, extracellular matrix, CD40 ligand, viral gene products neutral amino acids, zinc, estrogen and androgens. In contrast, stimuli which promote apoptosis include growth factors such as tumor necrosis factor (TNF), Fas, and transforming growth factor β (TGF β), neurotransmitters, growth factor withdrawal, loss of extracellular matrix attachment, intracellular calcium and glucocorticoids, for example. Other stimuli, including those of environmental and pathogenetic origins, also exist which can either induce or inhibit programmed cell death. Thus, apoptosis is mediated by diverse signals and complex interactions of cellular gene products which ultimately result in a cell death pathway that is required to regulate normal tissue homeostasis.

Several gene products which modulate the apoptotic process have now been identified. Although these products can in general be separated into two basic categories, gene products from each category can function to either inhibit or induce programmed cell death. One family of gene products are those which are members of the Bcl-2 family of proteins. Bcl-2, is the best characterized member of this family and inhibits apoptosis when overexpressed in cells. Other members of

this gene family include, for example, Bax, Bak, Bcl-x_L, Bcl-x_S, and Bad. While some of these proteins can prevent apoptosis, others such as Bax, Bcl-x_S and Bak function to augment apoptosis.

5 A second family of gene products which modulate
the apoptotic process is the family of proteases known as
aspartate-specific cysteine proteases (ASCPs). These
proteases are related genetically to the ced-3 gene
product which was initially shown to be required for
10 programmed cell death in the roundworm, *C. elegans*. The
ASCPs family of proteases includes, for example, human
ICE (interleukin-1- β converting enzyme), ICH-1_L, ICH-1_S,
CPP32, Mch2, Mch3, ICH-2 and ICE_{rel}-III. One common
feature of these gene products is that they are cysteine
15 proteases with specificity for substrate cleavage at Asp-
X bonds. Although these proteases induce cell death
when expressed in cells, several alternative structural
forms such as ICE δ , ICE ϵ , ICH-1_S and Mch2 β are known which
actually function to inhibit apoptosis.

20 In addition to the Bcl-2 and ASCP gene families
which play a role in apoptosis in mammalian cells, it has
become increasingly apparent that other gene products
exist which are important in mammalian cell death and
which have yet to be identified. For example, in
25 addition to Ced-3, another *C. elegans* gene known as Ced-4
exists which is also required for programmed cell death
in *C. elegans*. However, mammalian homologues of this
protein remain elusive and have not yet been identified.
Further, it is ambiguous as to whether other genes exist

use human proteins for the production of biopharmaceuticals as well as other molecules or compounds which are produced to modify the function of a regulator of programmed cell death. The use of human proteins as both biopharmaceuticals and to identify compounds is also important in regard to the therapeutic efficacy and safety of the treatment. For example, the half-life of non-human proteins can be comprised in a heterologous environment and the induction of a host immune response against a non-human protein can be potentially dangerous to the patient.

Thus, there exists a need to identify new apoptotic genes and their human homologues and for methods of modulating apoptotic process for the therapeutic treatment of human diseases. The present invention satisfies this need and provides related advantages as well.

SUMMARY OF THE INVENTION

The invention provides an isolated gene and an isolated nucleic acid sequence encoding human Bad and functional fragments thereof. Also provided is an isolated human Bad polypeptide and functional fragments thereof. Methods of identifying human Bad binding partners and methods of screening for compounds which interfere with the association of human Bad interacting polypeptides with human Bad are also provided. Finally, methods for decreasing or increasing the viability of a cell are provided as well.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide and predicted amino acid sequence of human Bad (SEQ ID NOS:1 and 2, respectively).

5 Figure 2 shows a comparison of the amino acid sequence between human Bad (SEQ ID NO: 2; clone PBM1) and mouse Bad (SEQ ID NO: 3; PmBad). The boxed regions emphasize amino acid identities and differences.

10 Figure 3 shows binding interactions between human Bad and Bcl-2 and between human Bad and Bcl-X_L in a yeast two-hybrid assay.

 Figure 4 shows the binding interactions between human Bad and Bcl-2 and between human Bad and Bcl-X_L in an ELIZA assay.

15

DETAILED DESCRIPTION OF THE INVENTION

 This invention is directed to the human Bcl-x_L/Bcl-2 associated death (Bad) promoting polypeptide and encoding nucleic acids. This protein is a member of the Bcl-2 family of cell death regulatory proteins in that it shows sequence homology with this family of proteins and is able to bind Bcl-x_L and to Bcl-2. The binding of human Bad to Bcl-x_L results in the induction of programmed cell death. The binding properties of human Bad to Bcl-x_L have been found to be different from the properties of Bax and Bak, two other death-enhancing Bcl-2 family proteins.

20

25

For example, amino-terminal deletions in Bax which create a fragment leaving the BH-1 and BH-2 domains intact completely abolish the ability of this fragment to interact with Bcl-X_L. In contrast, equivalent truncations of Bad leave the ability of this fragment to bind Bcl-x_L intact. This difference in interaction can have therapeutic advantages because compounds can be used which selectively target interactions of Bad and suppress programmed cell death without interfering with other cell death regulatory proteins.

In one embodiment, the invention is directed to nucleic acids encoding human Bad. The nucleic acids are used to produce recombinant Bad polypeptides which can be used in screening assays to identify compounds that interfere with Bad function. These screening assays yield compounds that antagonize the function of Bad and thus will prevent apoptosis. Such pharmaceutical compounds are useful for the treatment or prevention of diseases which are characterized by apoptotic cell death. Alternatively, assays can be configured to yield compounds that exhibit substantially the same function as, or "mimic" the function of Bad and thus enhance apoptosis. These pharmaceutical compounds are useful for the treatment or prevention of diseases which are characterized by the loss of apoptotic cell death. Human Bad polypeptides can also be used to identify novel interacting polypeptides and to identify the binding domains within these proteins as well as within known binding polypeptides such as Bcl-X_L. Similarly, these novel binding polypeptides and binding domains can be

used to inhibit or enhance apoptosis or to identify compounds which modulate their apoptotic function.

As used herein, the term "substantially" when referring to the human Bad nucleotide or amino acid sequence is intended to refer to the degree to which two sequences of between about 15-30 or more nucleotides in length, are identical or similar so as to be considered by those skilled in the art to be functionally equivalent. For example, the human Bad nucleic acid of the invention has a nucleotide sequence substantially the same as that shown in Figure 1 and as SEQ ID NO:1. Thus, if a second sequence is substantially the same as that shown as SEQ ID NO:1 so as to selectively hybridize, then it is considered functionally equivalent by those skilled in the art. Methods for sequence comparisons and determinations of similarity are well known and routine within the art.

Functionally equivalent nucleic acid sequences include, for example, sequences that are related, but different and encode the same human Bad polypeptide due to the degeneracy of the genetic code as well as sequences that are related, but different and encode a different human Bad polypeptide that exhibits similar functional activity. In both cases, the nucleic acids encode functionally equivalent gene products. Functional fragments of human Bad encoding nucleic acids such as oligonucleotides, polyoligonucleotides, primers and modified forms thereof are also considered to be within the definition of the term and the invention as claimed.

Functional equivalency is also relevant to human Bad

5 nucleic acids which do not encode gene products, for example, but instead are functional elements in and of themselves. Specific examples of such functional nucleic acids include, for example, promoters, enhances and other gene expression regulatory elements.

Human Bad polypeptides of the invention have an amino acid sequence substantially similar to that shown in Figures 1 and in SEQ ID NO:2. Functionally equivalent human Bad amino acid sequences similarly includes, for example, related, but different sequences so long as the different polypeptide exhibits at least one functional activity of human Bad. Specific examples of Bad polypeptide functional activities include functions such as the ability to bind or otherwise interact with Bad interacting polypeptides and/or Bad binding partners and the ability to promote cell death by heterodimerization with, for example, Bcl-x_l and Bcl-2. Related, but different polypeptides also include, for example, substitutions of conserved and/or non-essential amino acids within the Bad polypeptide sequence. Fragments and functional domains of human Bad are similarly included within the definition of the term and the claimed invention.

Therefore, it is understood that limited modifications may be made without destroying the

biological function of the human Bad polypeptide and that only a portion of the entire primary structure may be required in order to effect activity. For example, minor modifications of the human Bad amino acid sequence (SEQ
5 ID NO:2) which does not destroy its activity also fall within the definition of human Bad and within the definition of the polypeptide claimed as such. Also, for example, genetically engineered fragments of human Bad either alone or fused to heterologous proteins such as
10 fusion proteins that retain measurable enzymatic or other biological activity fall within the definition of the polypeptides as claimed.

It is understood that minor modifications of primary amino acid sequence may result in polypeptides
15 which have substantially equivalent or enhanced function as compared to the sequences set forth in Figure 1 (SEQ ID NO:2). These modifications may be deliberate, as through site-directed mutagenesis, or may be accidental such as through mutation in hosts which are human Bad
20 producers. All of these modifications are included as long as human Bad biological function is retained. Further, various molecules can be attached to human Bad, including for example, other proteins, carbohydrates, lipids, or chemical moieties. Such modifications are
25 included within the definition of human Bad polypeptides.

As used herein, the term "Bad interacting polypeptide" is intended to mean a peptide, polypeptide or protein which interacts directly or indirectly with the human Bad polypeptide. Such polypeptides include,

for example, Bcl-x_L and Bcl-2. When associated with human Bad, these interacting polypeptides form a heterodimer with the human Bad polypeptide. Direct interaction can include the direct binding of human Bad polypeptide with the interacting polypeptide. Binding can occur, for example, through polypeptide-polypeptide binding interaction or through binding involving post-translational modifications. Examples of post-translational modifications can include phosphorylation, glycosylation, acylation and lipidation. Indirect interaction between Bad and Bad interacting polypeptides can include, for example, the interaction of human Bad with a docking protein such as an SH2 or SH3 containing protein with functions to regulate the activity of a cell death protein proximal in the human Bad pathway. Human Bad interacting polypeptides can enhance Bad activity such as by augmenting the interaction of Bad with Bcl-x_L or Bcl-2 or with other cell death regulatory molecules. Alternative, human Bad interacting polypeptides can inhibit Bad activity by preventing the interaction of Bad with, for example, Bcl-x_L, Bcl-2 or other cell death regulatory molecules. Similarly, such interactions can occur directly or indirectly.

As used herein, the term "human Bad binding partner" is intended to mean polypeptides, non-polypeptide mimetics, organic molecules and other compounds which bind to the human Bad polypeptide. As with human Bad interacting polypeptide, human Bad binding partners can either inhibit or enhance the function of Bad through the binding of these partners to human Bad.

The invention provides a gene encoding human Bad or functional fragment thereof. The invention also provides an isolated nucleic acid sequence encoding human Bad or fragment thereof. The gene and nucleic acid sequences encode substantially the sequence as shown in SEQ ID NO:1. Fragments of the gene or nucleic acid sequence are provided which comprise single or double stranded nucleic acids having substantially the sequences shown in SEQ ID NO:1.

The human Bad nucleic acids of the present invention were identified by searching a human database of expressed sequence tags (ESTs) to identify potential new sequence fragments which may have homology to the mouse Bad gene and other Bcl-2 family cell death regulators. This approach was chosen because the standard approach of using the mouse sequence to probe human cDNA libraries had failed in this case as described in the Examples. Searches of the EST database identified short ESTs containing partial nucleotide sequence homology to the mouse Bad nucleotide sequence. These EST sequences were then used to design probes for the screening of cDNA libraries. Screening with the EST derived probes resulted in the identification of a putative human Bad nucleotide sequence which encoded a protein with approximately 77% amino acid homology to mouse Bad with the exception of an apparent truncation of 42 amino acids at the amino terminus. A comparison of the amino acid sequence between human Bad (SEQ ID NO: 2) and mouse Bad (SEQ ID NO: 3) is shown in Figure 2.

0902378-080301

Unexpectedly, this apparent truncation has now been shown to be the authentic amino terminus of human Bad.

In regard to the search for potential new sequences having homology to mouse Bad and other Bcl-2 family members, there are numerous considerations which must be taken into account when searching for homologies from unknown sequences such as ESTs fragments. For example, searching a genetic data base will yield homologous sequence matches to essentially any query nucleotide sequence. However, in order for the search to result in a meaningful match, additional criteria and permutations must be used to identify the authentic homologue of interest from among the non-specific homology matches. Mouse Bad shares the highest degree of homology with other Bcl-2 related proteins in the region believed to contain the binding site with Bcl-X_L and Bcl-2. A given EST returned by the search may not include one of these functionally important highly homologous sites, but rather, may only include a region within the polypeptide with cryptic homology. Confirming an EST as an actual Bad or BH1 and BH2 encoding polypeptide involves translation of all positive EST hits in three different reading frames and subsequent identification of conservative sequence homologies to the gene of interest. Then, using conventional cDNA cloning, a full length cDNA of the putative novel gene can be obtained and 1) analyzed for overall structural homology to Bad and Bcl-2 family members, 2) recombinantly expressed and analyzed for binding activity to other Bcl-2 family proteins and 3) analyzed for the induction

or suppression of programmed cell death by heterologous expression of the cDNA in appropriate cells.

Alternative methods than that described above for isolating human Bad encoding nucleic acids can similarly be employed. For example, using the teachings described herein, those skilled in the art can routinely isolate and manipulate human Bad nucleic acids using methods well known in the art. All that is necessary is the sequence of the human Bad encoding nucleic acids (Figures 1 and SEQ ID NO:1) or its amino acid sequence (Figures 1 and SEQ ID NO:2). Such methods include, for example, screening a cDNA or genomic library by using synthetic oligonucleotides, nucleic acid fragments or primers as hybridization probes. Alternatively, antibodies to the human Bad amino acid sequence or fragments thereof can be generated and used to screen an expression library to isolate human Bad encoding nucleic acids. Other binding reagents to human Bad polypeptides, such as Bcl-X_L or fragments thereof can similarly be used to isolate human Bad polypeptides having substantially the amino acid sequence show in Figure 1.

In addition, recombinant DNA methods currently used by those skilled in the art include the polymerase chain reaction (PCR) which, combined with the human Bad nucleotide and amino acid sequences described herein, allows reproduction of human Bad encoding sequences. Desired sequences can be amplified exponentially starting from as little as a single gene copy by means of PCR. The PCR technology is the subject matter of United States

Patent Nos. 4,683,195, 4,800,159, 4,754,065, and 4,683,202 all of which are incorporated by reference herein.

The above described methods are known to those skilled in the art and are described, for example, in Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, New York (1992) and the various references cited therein and in Ansubel et al., Current Protocols in Molecular Biology, John Wiley and Sons, Baltimore, MD (1989); and in Harlow et al., Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, New York (1989). These references and the publications cited therein are hereby expressly incorporated herein by reference.

The invention provides an isolated human Bad polypeptide comprising substantially the amino acid sequence as that shown in Figure 1 (SEQ ID NO:2). Human Bad functional fragments are also provided. A specific example of a human Bad functional fragment is the Bcl-X_L binding domain.

Isolated human Bad polypeptides of the invention can be obtained by a variety of methods known within the art. For example, the isolated peptides can be purified by biochemical methods including, for example, affinity chromatography. Affinity matrices which can be used for human Bad isolation can be anti-human Bad monoclonal or polyclonal antibodies prepared against the amino acid sequence shown in Figure

1 (SEQ ID NO:2), or fragments thereof such as synthetic peptides, recombinant fragments or the like.

Alternatively, cognate binding domains or polypeptides as well as other compounds known in the art which

- 5 specifically bind to human Bad can similarly be used as affinity matrices to isolate substantially pure human Bad polypeptides of the invention.

Human Bad polypeptides can also be produced by recombinant methods known to those skilled in the art.

- 10 Recombinant human Bad polypeptides include, for example, an amino acid sequence substantially the same as that shown in Figures 1 (SEQ ID NO:2) as well as fusion proteins and fragments thereof. The human Bad encoding nucleic acids can be cloned into the appropriate vectors
- 15 for propagation, manipulation and expression. Such vectors are known or can be constructed by those skilled in the art and should contain all expression elements necessary for the transcription, translation, regulation, and if desired, sorting of the human Bad polypeptides.
- 20 The vectors can also be for use in either prokaryotic or eukaryotic host systems so long as the expression and regulatory elements are of compatible origin. One of ordinary skill in the art will know which host systems are compatible with a particular vector. The recombinant
- 25 polypeptides produced can be isolated by, for example, the methods described above.

Apoptosis plays a significant role in numerous pathological conditions in that programmed cell death is either inhibited, resulting in increased cell survival,

or enhanced which results in the loss of cell viability. Examples of pathological conditions in which increased cell survival contributes to the disease process include cancers in general, and more particularly cancers such as lymphomas and hormone dependent tumors. Such hormone dependent tumors include, for example, breast, prostate and ovarian cancer. Inhibition of programmed cell death of immune cells plays a role in autoimmune diseases such as systemic lupus erythematosus and immune-mediated glomerulonephritis. Viruses such as herpesvirus, poxvirus and adenovirus carry anti-apoptotic genes to prolong production of viral particles by the host cell by preventing the host cell from committing suicide.

In contrast, apoptotic diseases where enhanced programmed cell death is part of the disease process generally include, for example, neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, and Cerebellar degeneration. Other diseases associated with increased apoptosis include, for example, myelodysplastic syndromes such as aplastic anemia and ischemic injury including myocardial infarction, stroke and reperfusion injury.

The human Bad encoding nucleic acids and polypeptides of the invention can be used to diagnose, treat or reduce the severity of cell death mediated diseases such as those described above as well as other diseases mediated by either increased or decreased programmed cell death. Additionally, the human Bad

encoding nucleic acids and polypeptides of the invention can be used to screen for pharmaceutical compounds and macromolecules which inhibit or promote human Bad mediated programmed cell death.

5 For example, the human Bad encoding nucleic acids, polypeptides and functional fragments thereof can be used to diagnose, or to generate reagents to diagnose diseases mediated or characterized by programmed cell death. Diagnosis can be by, for example, nucleic acid
10 probe hybridization with human Bad containing nucleotide sequences or by antibody or ligand mediated detection with human Bad binding agents. Such methods are routine to those skilled in the art. Detection can be performed ex vivo, for example, by removing a cell or tissue sample
15 from an individual exhibiting or suspected of exhibiting a cell death mediated disease. Correlation of increased human Bad expression or activity is indicative of diseases characterized by enhanced programmed cell death whereas correlation of decreased human Bad expression or
20 activity is indicative of diseases characterized by the suppression of programmed cell death.

The above human Bad polypeptides can also be formulated into pharmaceutical compositions known within the art for the treatment of cell death mediated diseases
25 characterized by increased cell survival and proliferation. Functional fragments, including peptides and peptide mimetics corresponding to the Bcl-X_L binding domain of human Bad can similarly be formulated for the treatment of such diseases associated with increased cell

survival and proliferation. Administration of human Bad polypeptides and functional fragments thereof will induce apoptosis in treated cells and eliminate those cells characterized by increased cell survival or

5 proliferation. Administration of non-human Bad polypeptides, molecules or compounds that induce the activation, regulation or expression of human Bad can similarly be used for the treatment of diseases characterized by increased cell survival and

10 proliferation. Specific examples of such polypeptides, molecules and compounds include Bad interacting polypeptides, binding partners and non-peptide compound equivalents.

To be effective, the human Bad polypeptides

15 must be introduced into the cells characterized by increased cell survival. Introduction can be accomplished by a variety of means known within the art including, for example, lipid vesicles and receptor mediated endocytosis. Targeting to the appropriate cell

20 type can similarly be accomplished through conjugation to specific receptor ligands, specific target cell antibodies and the like.

The human Bad polypeptides are administered by conventional methods, in dosages which are sufficient to

25 induce apoptosis in the cells characterized by increased cell survival or proliferation. Such dosages are known or can be easily determined by those skilled in the art. Administration can be accomplished by, for example, intravenous, intraperitoneal or subcutaneous injection.

Administration can be performed in a variety of different regimens which include single high dose administration or repeated small dose administration or a combination of both. The dosing will depend on the cell type,
5 progression of the disease and overall health of the individual and will be known or can be determined by those skilled in the art.

In contrast to the induction of human Bad mediated apoptosis for the treatment of pathological
10 conditions characterized by increased cell survival or proliferation, inhibitors of human Bad can be used to treat diseases characterized by increased programmed cell death. Such inhibitors can be, for example, binding domain inhibitors, including small non-peptide, organic
15 molecule inhibitors. Such inhibitors are formulated in a medium which allows introduction into the desired cell type. Alternatively, such inhibitors can be attached to targeting ligands for introduction by cell mediated endocytosis and other receptor mediated events.

20 Other inhibitors of human Bad include, for example, small molecules and organic compounds which bind and inactivate human Bad by a competitive or non-competitive type mechanism. Molecules, polypeptides or compounds which indirectly inhibit the human Bad pathway
25 can also be used as inhibitors of human Bad. Indirect inhibition can be accomplished by, for example, inhibition of Bad interacting polypeptides with Bad. Such inhibition includes, for example, the inhibition of binding events as well as post-translational

modifications which function in promoting human Bad mediated cell death. Human Bad inhibitors can be identified by screening for molecules which demonstrate specific or beneficial human Bad inhibitory activity.

- 5 Such methods are described further below and can be practiced by those skilled in the art given the human Bad nucleotide and amino acid sequences described herein.

- Treatment or reduction of the severity of cell death mediated diseases can also be accomplished by, for example, introducing expressible nucleic acids encoding human Bad polypeptides or functional fragments thereof into cells characterized by such diseases. For example, elevated synthesis rates of human Bad can be achieved by, for example, using recombinant expression vectors and gene transfer technology. Similarly, treatment or reduction of the severity of cell death mediated diseases can also be accomplished by introducing and expressing antisense human Bad nucleic acids so as to inhibit the synthesis rates of human Bad. Such methods are well known within the art and include, for example, the introduction of a compatible expression vector into population of cells characterized by either increased or decreased programmed cell death. Vectors can include, for example, both DNA and RNA viral vectors. Adenovirus is a specific example of a DNA viral vector.

Other methods of antisense inhibition known in the art include, for example, the use of synthetic oligonucleotides which are permeable to the cell membrane. Such oligonucleotides can include

methyolphosphonate and thiophosphonate chemical moieties for facilitating membrane permeability. The oligonucleotide can be employed for antisense inhibition of RNA processing, transport or translation events, or
5 alternatively, oligonucleotides which form triple helices can be used for the inhibition of transcription. Such methods are well known in the art.

The vector to be used in the methods of the invention will depend on desired cell type to be
10 targeted. For example, if neurodegenerative diseases are to be treated by decreasing the human Bad activity of affected neuronal cells then a vector specific for cells of the neuronal cell lineage should be used. Likewise, if diseases or pathological conditions of the
15 hematopoietic system are to be treated, than a viral vector that is specific for blood cells and their precursors, preferably for the specific type of hematopoietic cell, should be used. Moreover, such vectors can additionally be modified with specific
20 receptors or ligands and the like to modify or alter target specificity through receptor mediated events. These modification procedures can be performed by, for example, recombinant DNA techniques or synthetic chemistry procedures. The specific type of vector will
25 depend upon the intended application. The actual vectors are also known and readily available within the art or can be constructed by one skilled in the art using methodology well known to those skilled in the art.

Vectors containing human Bad nucleic acids or inhibitors of human Bad such as antisense nucleic acids can be administered in several ways to obtain expression and therefore either increase or decrease the activity of human Bad in the cells affected by the disease or pathological condition. If viral vectors are used, for example, the procedure can take advantage of their target specificity and consequently, do not have to be administered locally at the diseased site. However, local administration can provide a quicker and more effective treatment. Administration can also be performed by, for example, intravenous or subcutaneous injection into the subject. Injection of the viral vectors into the spinal fluid can also be used as a mode of administration, especially in the case of neurodegenerative diseases.

Transfection vehicles such as liposomes can be used to introduce the non-viral vectors into recipient cells within and inoculated area. Such transfection vehicles are known by one skilled within the art. Alternatively, non-targeting vectors can be administered directly into a tissue of any individual. Such methods are known within the art and are described by, for example, Wolff et al. (Science 247:1465-1468 (1990)).

Additional features can be added to the vectors to ensure safety and/or enhance therapeutic efficacy. Such features include, for example, markers that can be used to negatively select against cells infected with the recombinant virus. An example of such a negative

selection marker is the TK gene described above that confers sensitivity to the antibiotic gancyclovir. Negative selection is therefore a means by which infection can be controlled because it provides inducible suicide through the addition of antibiotic. Such protection ensures that if, for example, mutations arise that produce mutant forms of human Bad, dysfunction of apoptosis will not occur.

As described previously, the human Bad encoding nucleic acids and human Bad polypeptides of the invention can be used to screen for compounds which inhibit or enhance the activity of human Bad. Such compounds can act directly or indirectly on human Bad, human Bad binding partners or human Bad interacting polypeptides. Such screening methods are known to those skilled in the art and can be performed by either *in vitro* or *in vivo* procedures. For example, described in Example II is a specific *in vivo* assay for human Bad binding activity. Also described therein is the identification of a human Bad binding domain which can be used as an inhibitory compound for human Bad binding activity. Moreover, this binding domain can also be used to model alternative inhibitors of human Bad and human Bad binding partners. Methods other than that described in Example II can also be used to screen and identify compounds which inhibit human Bad. Specific examples include the *in vitro* screening of small molecule libraries such as those produced by combinatorial chemistry. Phage display peptide libraries can similarly be screened for inhibitory polypeptides. Such libraries offer the

advantages of producing greater than 10^8 peptide sequences which can be screened in a single round of panning. These methods as well as others are known within the art and can be utilized to identify compounds which inhibit or enhance human Bad binding activity.

It is understood that modifications which do not substantially affect the activity of the various embodiments of this invention are also included within the definition of the invention provided herein.

10 Accordingly, the following examples are intended to illustrate but not limit the present invention.

EXAMPLE I

Cloning And Characterization of Human Bad

This Example shows the cloning and sequence analysis of human Bad. The results described herein indicate that human Bad is a unique member of the BH1 and BH2 containing cell death regulatory polypeptides.

For the isolation of the human bad cDNA, an EST database was searched using the mouse protein sequence as a search criteria. However, prior to the use of the below described approach using expressed sequence tags (EST), two human c-DNA libraries were initially screened with a mouse bad probe under low stringency conditions. No human bad clones were identified using this approach.

25 An EST derived probe subsequently led to the isolation of human bad c-DNA clones from the same libraries.

Subsequent to the above conventional approach, the EST database (NCBI) was searched with a 204 amino acid mouse protein sequence using the BLAST network service. Initial search attempts resulted in partial sequences showing limited homology to the search query. Briefly, an EST sequence was identified that had 25 out of 32 identical amino acids in the region corresponding to mouse residues 100-131 and 7 out of 7 identical amino acids in a region corresponding to mouse residues 147-153. Overall, this EST sequence was 364 nucleotides in length and exhibited an overall sequence homology to the mouse sequence of 72% at the nucleotide level. Differences in nucleotide sequence between the actual EST and the corresponding mouse sequence were apparent and likely resulted from no less than 37 sequencing errors (additions, deletions and ambiguities) causing amino acid frame shifts. As a result, only 32 out of a potential 109 amino acids (29%) actually aligned with the query sequence. Nevertheless, this partial EST sequence homology tentatively identifies a partial cDNA encoding human Bad (EST #171560, GenBank #H18135) and was used to screen human cDNA libraries so as to identify, characterize and confirm the existence of the human Bad cDNA and polypeptide.

Human cDNA libraries were screened using a 200 bp fragment amplified from the above putative human Bad EST. The 200 bp fragment was labeled with ³²P-dCTP and then used to screen libraries derived from human substantia nigra, bone marrow and placenta. Screening methods were used were those known in the art and which

are essentially described in Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, New York (1992). Briefly, 10^6 plaque forming units were screened per library using a commercially available hybridization buffer, Quick Hybe (Stratagene) containing the above described putative human Bad EST probe. To identify positive plaques, duplicate filters were washed three times in 1X SSC and exposed to X-OMAT film (Kodak).

10 Screening of the each library resulted in the identification of putative cDNAs encoding human Bad. Surprisingly, none of these cDNAs appeared to be full length when compared to the corresponding mouse Bad sequence. For example, two partial cDNA's were isolated
15 from substantia nigra and encoded 95 amino acids of the 3' end of Bad. Similarly, seven partial cDNAs were isolated from a placental cDNA library and three from a bone marrow library. Again, these cDNAs were truncated in that they encoded sequences substantially shorter than
20 the mouse Bad sequence. A sequence comparison of each of these cDNAs revealed that all shared identical nucleotide sequence in the region identified as the open reading frame. However, when compared to the mouse Bad nucleotide and predicted amino acid sequence all of these
25 isolated clones appeared to be truncated at their 5' ends.

Further sequence analysis of each of these clones revealed that the longest clone, termed BM1, exhibited a truncation of approximately 42 amino acids

compared to the initiation methionine of the mouse sequence. Nevertheless, these apparently partial human Bad cDNA clones exhibited a predicted open reading frame that initiated with an amino terminal methionine. A
5 further comparison of the nucleotide and encoding amino sequence revealed a 84% nucleotide identity and a 77% amino acid identity when compared to the mouse sequence. Thus, the similarity in sequence identity together with the apparent amino terminal truncation lead to the
10 initial conclusion that the longest of the human Bad cDNA clones was probably less than full length. Surprisingly however, after further attempts to isolate a longer clone and after numerous experimentations that independently assessed the apparent authenticity of the methionine as a
15 translation initiator present in human clones, it was concluded that the BM1 clone (SEQ ID NO:1) and shown in Figure 1 as human Bad was the full length human sequence. The predicted amino acid sequence of human Bad is also shown in Figure 1 and as SEQ ID NO:2. Figure 2 shows the
20 amino acid alignments of human and mouse Bad. The experimentation yielding these conclusions are described further below.

To verify that cDNA BM1 encodes full length human Bad, the 5' sequence of a number of different cDNAs
25 were analyzed. The results of that analysis revealed that seven different human placenta clones and three different human bone marrow clones terminated less than 54 nucleotides upstream from the above described proposed start methionine. In addition, two subsequently entered
30 EST sequences of 317 and 300 nucleotides in length and

having some similarity to the above described human Bad cDNA were also found to terminate in this region. In all cases, the sequence around the start methionine satisfied Kozac criteria (A at -3 position). Also isolated were two clones from placenta (PL14 and PL21) which differed from other isolates in sequence just 5' to above described start methionine. In contrast, the sequence in this 5' region contains a consensus site for an RNA splicing junction and is similar to the mouse Bad sequence. Taken together, the above results indicate that there can be alternatively spliced forms of human Bad.

EXAMPLE II

Human Bad Binding Partners

This Example shows that Human Bad can bind to Bcl-X_L.

To assess whether Bcl-2 family members bind to or otherwise interact with human Bad and to map their binding domains, a yeast two-hybrid system was employed using the above described clones. Briefly, the full length human Bad sequence was subcloned into the yeast 2-hybrid vector pEGE202, or "bait-vector" to express Bad as a fusion protein with the DNA-binding protein LexA (Ruden et al., Nature (1991) 350:426-430). Human Bcl-2, Bcl-X, Bax, Bak and Bad were similarly cloned into the yeast 2-hybrid vector pJG4-5, or "prey-vector" to expressed these Bcl-2 family members as fusion proteins with the transcriptional activator B42 (Gyuris et al., Cell (1993) 75:791-803). Briefly, the full length human

Bad sequence was amplified by PCR and subcloned into the bait-vector. For the PCR reaction we used the following primers: 5'ATC AGT GAA TTC ACT ATG TTC CAG ATC CCA GAC3' (SEQ ID NO: 4) and 5'ATC GAT CTC GAG TCA CTG GGA GGG GGC GGA GCT3' (SEQ ID NO: 5). The resultant PCR fragment was digested with EcoRI and Xho I and then subcloned into the bait vector. The human Bad sequence was also PCR amplified and subcloned into the prey-vector using the same primers and subcloning strategy as described above.

Human Bak was PCR amplified and subcloned as an EcoRI/XhoI fragment into vector pJG4-5 using the following primers: 5'ATC AGT GAA TTC ACT ATG GCT TCG GGG CAA GGC CC-3' (SEQ ID NO: 6) and 5' ATC GAT CTC GAG TCA GTT CAG GAT GGG ACC ATT GC-3 (SEQ ID NO: 7). Human Bax was PCR amplified and subcloned as an EcoRI/XhoI fragment into vector pJG4-5 using the following primers: 5' ATC AGT GAA TTC ACT ATG GAC GGG TCC GGG GAG-3' (SEQ ID NO: 8) and 5'-TAC AGT CTC GAG TCA GGT CAC GGT CTG CCA CGT GGG-3 (SEQ ID NO: 9). Human Bcl-2, Bcl-X-L and Bcl-X-S in the yeast 2-hybrid vector pJG4-5 were available and used as described previously by Sato et al. (Proc. Natl. Acad. Sci. 91:9238-9242 (1994)).

To determine if human Bad binds to any of the above Bcl-2 family members, the Bad-bait plasmid was cotransformed with every prey plasmid and the reporter plasmid p18-34 into the yeast strain EGY191. Cells were induced for 6 hours with galactose to allow expression from the prey plasmid. Expression was assayed by lysing a 100 ul aliquot of yeast cells (OD 600=0.6) in 96 well plates in Z-buffer (60 mM Na₂HPO₄, 40 mM NaH₂PO₄ x H₂O, 10

mM KCL, 1 mM MgSO₄ x 7 H₂O) and 1 u Lyticase (Boehringer Mannheim) for 1 hour followed by addition of the substrate ONPG into a final concentration of 2 mg/ml (Sigma, St. Louis, MO). The reaction was stopped with
 5 0.4 M NaOH and the plate was read at an OD 420 nm. Protein-protein interaction was measured as beta-galactosidase activity using the following equation:
 OD 420: (OD 600 x 0.1 ml x ONPG reaction time).

Results of these experiments showed that human
 10 Bad interacts with Bcl-2 and Bcl-X_L. However, no interactions were observed between human Bad and Bax, Bak or itself. These results are presented in Figure 3.

EXAMPLE III

In Vitro Screening and Identification of Compounds which 15 Modify the Binding Activity of Human Bad

This Example shows the screening and identification of compounds which inhibit human Bad interacting polypeptides and human Bad associations using recombinant human Bad polypeptides.

20 To screen for compounds which disrupt associations between human Bad interacting polypeptides and human Bad, an ELISA-type assay is employed. This assay format is used here due to its reliability and simplicity, however, other formats and binding assays can
 25 be substituted in place of the ELISA. In particular, the yeast-2-hybrid interaction between Bad and Bcl-X_L or Bcl-2

shown in Example II can also be used as a compound screening assay.

To generate fusion proteins for these screening assays, Bad and Bcl-xL were cloned into pET-15b (Novagen) and pGEX 4T1 (Pharmacia) vectors, respectively. The pET15b-bad construct was made as follows. Full length bad cDNA was amplified by PCR using cDNA BM1 as template and primers containing NdeI (5') and XhoI (3') restriction sites (5' primer: 5'-GGG AAT TCC ATA TGT TCC AGA TCC CAG AG-3' (SEQ ID NO: 10)); 3' primer: 5'-TAC AGT CTC GAG TCA CTG GGA GGG GGC GGA GCT-3' (SEQ ID NO: 11)). The PCR product was then digested with NdeI/XhoI and ligated into the NdeI and XhoI sites of pET15b. The above primer sequences were chosen such that the bad ORF sequence would be in frame with 5' vector sequence encoding a six residue Histidine tag. The final construct was termed pET-15 6H Bad.

The pGEX-Bcl-X_L and pGEX Bax were constructed in similar fashion using primers containing EcoRI sites on both 5' and 3' ends (Bcl-X_L, 5' primer: 5'-AGT ATC GAA TTC ATG TCT CAG AGC AAC CGG-3' (SEQ ID NO: 12), Bcl-X_L 3' primer: 5'-ATT GAT GAA TTC GTT GAA GCG TTC CTG GCC CTT-3' (SEQ ID NO: 13), Bax-5' primer: 5'-ATC AGT CTC GAG ACT ATG GAC GGG TCC GGG GAG-3' (SEQ ID NO: 14), Bax-3' primer: 5'-TAC GAT GAA TTC GGT CAC GGT CTG CCA CGT GGG-3' (SEQ ID NO: 15). PCR products were then digested with EcoRI and ligated into the EcoRI site of pGEX 4T1. The above primer sequences were chosen such

0922338-080994

that Bcl-X_L and Bax ORF sequence would be in frame with 5' vector sequence encoding a glutathione-S-transferase tag.

Proteins were purified according to methods known in the art and to protocols provided by the vendor. Briefly, pET 15-6Hbad was transformed into E. coli strain BL21 plysS and pGE constructs into strain XL1-Blue for protein production. Bacteria were grown in the presence of 100 µg/ml carbenicillin (and 34 µg/ml chloramphenicol in the case of BL21 plysS) until reaching an O.D of 1. The cultures were induced with 1 mM IPTG and grown for an additional 3 hours. Bacteria were harvested by centrifugation, and protein isolated by sonication in 50 mM Tris, pH7 with 0.1% Triton X-100 containing protease inhibitors PMSF, leupeptin (1 mg/ml) and benzamidine (1 mM). Proteins were then purified by affinity chromatography using either nickel (for 6HBad) or glutathione (for GST Bcl-X_L and GST-Bax) as the affinity ligand.

To perform the screening assays, multiwell plates were coated overnight at 4°C with 50 µl/well of the 6HBad fusion protein (4µg/ml) in phosphate buffered saline (PBS). The plates were then washed twice with phosphate-buffered saline containing 0.02% Tween (PBST) and blocked with PBS containing 1% BSA (150 µl/well) for 2 hours at room temperature. Prior to addition of compounds to be tested, the plates were washed twice with PBST. Test compounds were added to the wells to a final concentration of about 20µM (50 µl/well) in PBST + 0.5% BSA. The plates were then incubated for 1 hour at room

5

10

15

20

25

5 Although the invention has been described with
reference to the disclosed embodiments, those skilled in
the art will readily appreciate that the specific
experiments detailed are only illustrative of the
invention. It should be understood that various
10 modifications can be made without departing from the
spirit of the invention. Accordingly, the invention is
limited only by the following claims.